

Abstrak

FAKTOR-FAKTOR YANG BERHUBUNGAN DENGAN KEJADIAN KANKER SERVIKS DI RSUD PROF. DR. MARGONO SOEKARJO PURWOKERTO

Silvia Salwa¹, Yunita Sari², Agis Taufik²

Latar Belakang: Kanker serviks menjadi penyakit pembunuh wanita nomer satu di Indonesia. Hal ini dikarenakan kurangnya informasi dan deteksi dini. Oleh karena itu, perlu adanya pencegahan yaitu mengetahui faktor apa saja yang berhubungan dengan kejadian kanker serviks.

Tujuan: Penelitian ini bertujuan untuk mengetahui faktor apa saja yang berhubungan dengan kejadian kanker serviks di RSUD Prof. dr. Margono Soekarjo Purwokerto.

Metode: Penelitian ini merupakan penelitian analitik dengan rancangan *cross sectional*. Teknik pengambilan sampel menggunakan *consecutive sampling* dengan jumlah sampel 50 responden di RSUD Prof. dr. Margono Soekarjo Purwokerto. Instrumen yang digunakan pada penelitian ini yaitu kuisioner. Data dianalisis secara univariat, bivariat, dan multivariat.

Hasil: Faktor yang berhubungan dengan kejadian kanker serviks yaitu *external vaginal douching* dengan p 0,021, usia pertama kali hamil dengan p 0,007, paritas dengan p 0,015, dan jarak persalinan dengan p 0,015. Faktor yang tidak berhubungan dengan kejadian kanker serviks yaitu jenis persalinan dengan p 1,000 dan riwayat abortus dengan p 0,442.

Kesimpulan: Faktor *external vaginal douching* menjadi faktor dominan yang berhubungan dengan kejadian kanker serviks dengan nilai OR sebesar 18,602 yang artinya bahwa *external vaginal douching* berisiko sebesar 18,602 kali terhadap kejadian kanker serviks.

Kata Kunci: Kanker Serviks, *External Vaginal Douching*, Usia Pertama Kali Hamil, Paritas, Jarak Persalinan

¹Mahasiswa Jurusan Keperawatan Fakultas Ilmu-Ilmu Kesehatan Universitas Jenderal Soedirman

²Departemen Keperawatan Medikal Bedah Jurusan Keperawatan Fakultas Ilmu-Ilmu Kesehatan Universitas Jenderal Soedirman

Abstract

FACTORS ASSOCIATED WITH CERVICAL CANCER IN RSUD PROF. DR. MARGONO SOEKARJO PURWOKERTO

Silvia Salwa¹, Yunita Sari², Agis Taufik²

Background: Cervical cancer has become Indonesia's number one disease that cause death of females. This is due to insufficient information and early detection. Therefore, prevention is needed which that is to recognize the factors that can cause cervical cancer.

Purpose: This research was aimed to determine any factors that are related to cervical cancer at RSUD Prof. dr. Margono Soekarjo Purwokerto.

Methods: This research was an analytical research with a cross sectional design. The technique that was used to collect samples was a consecutive sampling with the total samples of 50 responders at RSUD Prof. dr. Margono Soekarjo Purwokerto. The instrument that was used in this research was a questionnaires. Data analysed univariate, bivariants, and multivariates.

Results: Factors that were related to cervical cancer are external vaginal douching factor with p 0.021, first stage in pregnancy with p 0.007, parity with p 0.015, and childbirth with p 0.015. A factor that is not related to the cervical cancer is abortus breeding with p 1.000 and abortive history with p 0.442.

Conclusion: The external vaginal douching factor has become the dominant factor that is associated with cervical cancer which means that the external vaginal douching is at 18.602 risk for cervical cancer.

Keywords: Cervical Cancer, External Vaginal Douching, First Stage In Pregnancy, Parity, Childbirth

¹Nursing Departement Student of Health Sciences Faculty of Jederal Soedirman University

²Departement of Medical Surgery Nursing of Nursing Departement of Health Sciences Faculty of Jendral Soedirman University